

BIBLIOGRAPHIE

INTRODUCTION ET CHAPITRE 1 : STOCKAGE DES DECHETS MENAGERS ET INDUSTRIELS

Ademe. Etude bibliographique des rejets des différentes techniques de traitement des résidus urbains.; 1995. Etude INSAVALOR POLDEN pour le compte de l'Ademe. 262p.

Ademe. Techniques de gestion des déchets ménagers - Etat des connaissances. Guide de révision des Plans Départementaux. 2000. Réf. Ademe 3567. 94p.

Agences de l'Eau, Ministère de l'Environnement. Etude inter-Agences de l'eau : Evaluation des flux polluants liquides générés par les décharges de déchets ménagers. 1997.

AGHTM. La décharge a un avenir : le centre de stockage. Dossier AGHTM. TSM 2000;1:5-54.

Amokrane A. Epuration des lixiviats de décharge. Prétraitement par coagulation-floculation. Traitement par osmose inverse. Post-traitement par incinération. Thèse INSA de Lyon (LCPAE); 1994. 254p.

Barres et al. Les lixiviats de décharges. Le point des connaissances en 1990. T.S.M.-L'eau 1990;85(6):289-313.

Borgès de Castilhos JA. Simulation au laboratoire du comportement hydraulique et de l'évolution bio-physico-chimique d'un déchet type composite mis en décharge: Thèse INSA de Lyon (LCPAE); 1988. 184p.

Bryant MP. Microbial methane production. Theoretical aspects. Journal of Animal Science 1979;48(1):193-201.

Chian ES, Pohland FG, Chang KC, Harper SR. Leachate generation and control at landfill disposal sites. In: Int. Conf. "New directions and research in waste treatment and residuals management"; juin 1985; Vancouver; p.745-752.

Christensen TH, Cossu R, Stegmann R. Gas issues in landfill management. In: Landfilling of waste : biogas. E&F Spon. Londres; 1996. p. 3-7.

Deloraine A, Zmirou D, Tillier C, Boucharlat A, Bouti H. Case-control assessment of the short-term health effects of an industrial toxic waste landfill. Environ.Res. 1995;68(2):124-132.

Dolk H, Vrijheid M, Armstrong B, Abramsky L, Bianchi F, Garne E et al. Risk of congenital anomalies near hazardous-waste landfill sites in Europe: the EUROHAZCON study. Lancet 1998;352(9126):423-427.

Dor F. Santé-Déchets: Certitudes et Incertitudes. Environnement et Technique 1998;176:27-29.

Farquhar GJ, Rovers FA. Gas production during refuse decomposition. Water, Air, Soil Pollution 1973;2:483-493.

Lombard A. Centres d'enfouissement technique : en route vers le zéro rejet. L'Eau, l'Industrie, les Nuisances 1999;227:33-35.

Mortgat B. Maîtrise du risque étanchéité des centres de stockage de déchets ménagers. *Environnement & Technique* 1997;171:29-33.

Pohland FG, Harper SR. Critical review and summary of leachate and gas production from landfills.: Georgia Institute of Technology (Atlanta, USA); 1985. Project n°E-20-G01, EPA CR 809 997. 165p.

Poulleau J. Caractérisation des biogaz. Bibliographie. Mesures sur site. Ineris; 2002. Rapport final. 31p. + annexes.

Rees JF. The fate of organic compounds in the landfill disposal of organic matter. *Journal of Chem. Tech. Biotechnol.* 1980;30:161-175.

Shusterman D. Critical review : the health significance of environmental odor pollution. *Arch.Environ.Health* 1992;47:76-87.

Touze-Foltz N. Détection, localisation et contrôle des fuites dans les géomembranes. *Environnement & Technique* 2003;223:21-25.

Zmirou D, Deloraine A, Saviuc P, Tillier C, Boucharlat A, Maury N. Short-term health effects of an industrial toxic waste landfill: a retrospective follow-up study in Montchanin, France. *Arch.Environ.Health* 1994;49(4):228-238.

CHAPITRE 2 CONNAISSANCES ACTUELLES

Abatemarco DJ, Delnevo CD, Rosen M, Weidner BL, Gotsch AR. Medical surveillance practices of blue collar and white collar hazardous waste workers. *J. occup. environ. Med.* 1995;37(5):578-582.

Adams KJ, Preston CD. Evidence for the effects of atmospheric pollution on bryophytes from national and local recording. In: Harding P.T., *Biological recording of changes in British wildlife*. ITE Symposium 26; 1992.

Ademe. Etude bibliographique des rejets des différentes techniques de traitement des résidus urbains. 1995. Etude INSAVALOR POLDEN pour le compte de l'Ademe. 262p.

Adeola FO. Environmental hazards, health and racial inequity in hazardous waste distribution. *Environ.and behavior* 1994;26 (1):99-126.

AFNOR. Qualité de l'air - Méthode de mesurage de l'odeur d'un effluent gazeux. Détermination du seuil de perception. Norme X43-101; 1986.

Allen MR, Braithwaite A, Hills CC. Trace organic compounds in landfill gas at seven U.K. waste disposal sites. *Environ. Sci. Technol.* 1997;31:1054-1061.

Allen MR, Braitwaithe A, Hills CC. Trace organic compounds in landfill gas at seven U.K. waste disposal sites. *Environ. Sci. Technol.* 1997;31(4):1054-1061.

Amler MR, Gibertini M, Lybarger JA, Hall A, Kakolewski K, Phifer BL et al. Selective approaches to basic neurobehavioral testing of children in environmental health studies.

Neurotoxicol. Teratol. 1996;18(4):429-434.

Ammann K, Herzig R, Liebendörfer L, Urech M. Multivariate correlation of deposition data of 8 different air pollutants to lichen data in a small town in Switzerland. In: Verlag B, editor. *Advances in Aerobiology*. Basel; 1987. p. 401-406.

Amokrane A. Epuration des lixiviats de décharge. Prétraitement par coagulation-floculation. Traitement par osmose inverse. Post-traitement par incinération.: Thèse INSA de Lyon (LCPAE); 1994. 254p.

Anon. Hazardous-waste sites: priority health conditions and research strategies - United States. *Morb. Mort. wkly. Rep.* 1992a;41(5):72-74.

Anon. Hazardous-waste sites: priority health conditions and research strategies - United States. *J. Am. med. Assoc.* 1992b;267(9):1180.

Apostoli P, Bartoli D, Alessio L, Buchet JP. Biological monitoring of occupational exposure to inorganic arsenic. *Occup. Environ. Med* 1999;56(12):825-832.

Arthur MA, Rubin G, Schneider RE, Weinstein LH. Uptake and accumulation of selenium by terrestrial plants growing on a coal fly ash landfill. Part 3. Forbs and grasses. *Environ. Toxicol. Chem.* 1992;11:1301-1306.

Assmuth T, Kalevi K. Concentrations and toxicological significance of trace organic compounds in municipal solid waste landfill gas. *Chemosphere* 1992;24(9):1207-1216.

Asta J, Erhardt W, Ferretti M, Fornasier F, Kirschbaum U, Nimis PL et al. Mapping lichen diversity as an indicator of environmental quality. In: Kluwer, editor. Nimis P.L., Scheidegger C. & Wolseley P.A.. *Monitoring with lichens - Monitoring lichens*. The Netherlands; 2002. p. 273-279.

Asta J, Gombert S, Benoit-Guyot JL, Seigle-Murandi F, Steiman R, Lafond M et al. Evaluation des nuisances et impacts liés à l'incinération d'ordures ménagères et assimilés: GRIDEC-Ademe; 1996. 141p., 15 annexes.

Asta J, Guillard E, Tissut M, Gaude T, Ravanel P. Heavy metal transfer from atmosphere to plants. *J. Phys. IV France* 2003;107:65-67.

Asta J, Keck G. Utilisation des lichens et des lapins en tant que bioindicateurs. Etude du CERED, point zéro; 1995. Non publié.

Asta J, Keck G. Suivi du site de stockage du CERED par les organismes " sentinelles " : lichens et lapins. Suivi à deux ans (décembre 1997); 1998. Non publié.

Asta J, Keck G. Suivi de la contamination chimique du site de stockage du CERED par les organismes " sentinelles " : lichens et lapins. Suivi à quatre ans; 2000. Non publié.

Barbier R. Une société au rendez-vous de ses déchets. L'internalisation des déchets comme figure de la dynamique du collectif. Thèse de doctorat en socio-économie de l'innovation: Ecole des Mines de Paris, Centre de Sociologie de l'Innovation; 1996.

Barthe Y. Comment parle-t-on des générations futures? Paris; 1997.

Bernardet-Van Staëvel E. De la monstrosité du Déchet Ultime. Dir Jean-Claude Beaune. In: Le déchet, le rebut, le rien. Seyssel, Champ Vallon: Coll. Milieux; 1999.

Bernardet-Van Staëvel E. L'impact des déchets sur la santé: approche anthropologique des représentations sociales des salariés et des riverains d'usines d'incinération d'ordures ménagères. Thèse de doctorat d'anthropologie: Ecole des Hautes Etudes en Sciences Sociales / Ademe; 2003.

Bernardet-Van Staëvel E, Navel D, Berthet C. Aide à la conception d'outils de communication sur les déchets industriels. Ph. 1: Etude sémantique et discours des acteurs sociaux: RECORD; 1995.

Berny P, Lachaux O, Mazallon M, Buronfosse T, Gillet C. Zebra mussels as indicators of fresh water contamination with lindane. Environ. Res. 2002;19:2614-2620.

Berry M, Bove F. Birth weight reduction associated with residence near a hazardous waste landfill. Environ. Health Perspect. 1997;105:856-861.

Billeret M, Asta J, Buronfosse T, Keck G, Berny P. Monitoring strategy for municipal landfill sites : can rats be used as indicators of pollutant availability and toxicity? soumis.

Billeret M, Collomb B, Buronfosse T, Berny P. In vivo assessment of the genotoxicity of PAH-contaminated soils in rats exposed by natural routes. J. polycyclic arom Compds 2000;20:275-289.

Blandin P. Bio-indicateurs et diagnostic des systèmes écologiques. Bulletin d'Ecologie 1986;17:215-307.

Blatrix C. Le maire, le commissaire enquêteur et leur "public". In: PUF, editor. CURAPP/CRAPS, La démocratie locale. Paris; 1998.

Bord RJ, O'Connor RE. Determinants of risk perceptions of hazardous waste site,. Risk anal. 1992;12(3):411-416.

Bouché M. Earthworm toxicological tests, hazard assessment and biomonitoring. In: Publications SPB, editor. Edwards, C.A., Neuhauser, E.F. *Earthworm in Waste and Environmental Management*. La Haye; 1988. p. 315-320.

Bradley AD, Cook DJ, Edwards JS, Johnston AG, Linforth RST, Taylor AJ. The control and measurement of landfill odours. In: Proceedings of Sardinia; 1-5 octobre 2001; Cagliari, Italie; 2001. p. 681-690.

Brandt CA, Rickard WH. Detection of metal contamination in wild asparagus near a waste disposal site. Environ. Monit. Assess. 1996;43(3):201-216.

Brandvold LA, Popp BR, Swartz SJ. Lead content of plants and soils from three abandoned smelter areas in and near Socorro, New Mexico. Environ. Geochem. Health 1996;18:1-4.

Brodo I. Transplant experiments with corticolous lichens using a new technique. Ecology 1961;42:838-841.

Brown P. Popular epidemiology and toxic waste contamination: lay and professional ways of knowing. J.Health soc.Behav. 1992;33:267-281.

Brown P. Race, class and environmental health: a review and systemization of literature. *Environ.Res.* 1995;69(1):15-30.

Bukowski JA, Goldstein MD, Korn LR, Rudakewyck M, Shepperly D, Gates D et al. Chromium exposure assessment of outdoor workers in Hudson county. *NJ. Sci. Total Environ* 1992;122(3):291-300.

Burkart R. Consensus oriented public relations as a solution to the landfill conflict. *Waste management research* 1994;12:223-232.

Callon M, Lascoumes P, Barthe Y. *Agir dans un monde incertain. Essai sur la démocratie technique.* Seuil, Paris; 2001.

Castle G. Hazardous waste facility siting in Minesota. *Air Waste* 1993;43:963-969.

CEC (Commission of the European Communities). *Landfill gas - From environment to energy:* CEC; 1992.

Chateauraynaud F, Torny D. *Les sombres précurseurs. Une sociologie pragmatique de l'alerte et du risque;* Maisons des sciences de l'homme, 1999.

Chiriack R. *Etude des émissions des COV par les Centres d'Enfouissement Techniques d'OM. Evaluation de l'impact environnemental.* Thèse en cours, à la demande de l'Ademe, UCB Lyon 1. *Rapport intermédiaire;* 2002. 92p.

Choudhary G. Human health perspectives on environmental exposure to benzidine : a review. *Chemosphere* 1996;32(2):267-291.

Christensen TH, Cossu R, Stegmann R. Gas issues in landfill management. In: E&FN Spon, editor. *Landfilling of waste : biogas.* Londres; 1996a. p. 3-7.

Christensen TH, Kjeldsen P, Lindhardt BO. Gas-generating processes in landfills. In: E&FN Spon, editor. *Landfilling of waste : biogas.* Londres; 1996b. p. 27-50.

Cislaghi C, Nimis PL. Lichens, air pollution and lung cancer. *Nature* 1997;387:463-464.

Clark DL, Bickham JW, Baker DL, Cowman DF. Environmental contaminants in Texas, USA, wetlands reptiles: evaluation using blood samples. *Environ. Toxicol. Chem.* 2000;19(9):2259-2265.

Cobb GP, Sands K, Waters M, Wixson BG, Dorward-King E. Accumulation of heavy metals by vegetables grown in mine wastes. *Environ. Toxicol. Chem.* 2000;19:600-607.

Cohen JT, Beck BD, Rudel R. Life years lost at hazardous waste sites : remediation worker fatalities vs. cancer deaths to nearby residents. *Risk Anal.* 1997;17(4):419-425.

Costello RJ, King MV. Protecting workers who clean up hazardous waste sites. *Am. ind. Hyg. Assoc. J.* 1982;43(1):12-17.

Croen LA, Shaw GM, Sanbonmatsu L, Selvin S, Buffler PA. Maternal residential proximity to hazardous waste sites and risk for selected congenital malformations. *Epidemiology* 1997;8:347-354.

Cuny D, Pignata ML, Kranner I, Beckett R. Biomarkers of pollution-induced oxidative stress and membrane damage in lichens. In: Kluwer, editor. Nimis P.L., Scheidegger C. & Wolseley P.A.. *Monitoring with lichens - Monitoring lichens*. The Netherlands; 2002. p. 97-110.

Czarnowski W, Krechniak J. Fluorides in hair and urine of children in the vicinity of a phosphate industry waste disposal site. *Fluoride* 1990;23:119-122.

Danse IHR, Garb LG, Moore RH. Blood lead surveys of communities in proximity to lead-containing mill tailings. *Am. ind. Hyg. Assoc. J.* 1995;56(4):384-393.

Dayal H, Gupta S, Trieff N, Maieron D, Reich D. Symptom clusters in a community with chronic exposure to chemicals in two Superfund sites. *Arch. environ. Contam. Toxicol.* 1995;50(2):108-111.

De Souza Bueno AM, De Bragança Pereira CA, Rabello-Gay MN. Environmental genotoxicity evaluation using cytogenetic end points in wild rodents. *Environ. Health Perspect.* 2000;108:1165-1169.

Deipser A, Poller T, Stegmann R. Emissions of volatile halogenated hydrocarbons from landfills. In: E&FN Spon, editor. *Landfilling of waste : biogas*. Londres; 1996. p. 59-71.

Deipser A, Stegmann R. The origin and fate of volatile trace components in municipal solid waste landfills. *Waste management & research* 1994;12(2):129-139.

Deloraine A, Zmirou D, Tillier C, Boucharlat A, Bouti H. Case-control assessment of the short-term health effects of an industrial toxic waste landfill. *Environ.Res.* 1995;68(2):124-132.

Deportes I, Benoit-Guyod JL, Zmirou D. Hazard to man and the environment posed by the use of urban waste compost: a review. *Sci. total Environ.* 1995;172:197-222.

Diaz-Barriga F, Santos MA, Yanez L, Cuellar JA, Ostrosky-Wegman P, Montero R et al. Biological monitoring of workers at a recently opened hazardous waste disposal site. *J. Expo. Anal. environ. Epidemiol.* 1993;3(supp1):63-71.

Dolk H, Vrijheid M, Armstrong B, Abramsky L, Bianchi F, Garne E et al. Risk of congenital anomalies near hazardous-waste landfill sites in Europe: the EUROHAZCON study. *Lancet* 1998;352(9126):423-427.

Domingo JL, Schuhmacher M, Granero S, de Kok Ham. Temporal variation of PCDD/F levels in environmental samples collected near an old municipal waste incinerator. *Environ. Monit. Assess.* 2001b;69(2):175-193.

Domingo JL, Schuhmacher M, Llobet JM, Müller L, Rivera J. PCDD/F concentrations in soil and vegetation in the vicinity of a municipal waste incinerator after a pronounced decrease in the emissions of PCDD/Fs from the facility. *Chemosphere* 2001a;43:217-226.

Domingo JL, Schuhmacher M, Müller L, Rivera J, Granero S, Llobet JM. Evaluating the environmental impact of an old municipal waste incinerator: PCDD/F levels in soil and vegetation samples. *J. hazard. mater. A* 2000;76:1-12.

Dudka S, Piotrowska M, Terelak H. Transfer of cadmium, lead and zinc from industrially contaminated soil to crop plants. *Environ. Pollut.* 1996;94(2):181-188.

Dunn JR, Taylor SM, Elliott SJ, Walter SD. Psychosocial effects of PCB contamination and remediation : the case of Smithville Ontario. *Soc.Scie.Med.* 1994;39(8):1093-1104.

Dunne MP, Burnett P, Lawton J, Raphael B. The health effects of chemical waste in an urban community. *Med.J.Aust.* 1990;152:592-597.

Echaubard M. Les animaux comme indicateurs biologiques de pollution. In: *Proceedings of Marqueurs biologiques de pollution*; 21-22 septembre 1995; Chinon; 1995. p. 335-358.

Ehrig HJ. Quality and quantity of sanitary landfill leachate. *Waste management & research* 1983;1:53-68.

Ehrig HJ. Water and element balances of landfills. In: Baccini, editor. *The Landfill - Reactor and Final Storage - Swiss Workshop on Land Disposal of Solid Wastes - Gerzensee, March 14-17,1988*; 1989. p. 83-115.

Eklund B, Anderson EP, Walker BL, Burrows DB. Characterization of landfill gas composition at the Fresh Kills municipal solid-waste landfill. *Environ. Sci. Technol.* 1998;32(15):2233-2237.

Elliott P, Briggs D, Morris S, de Hoogh C. Risk of adverse birth outcomes near landfill sites. *Br. med. J.* 2001a;323(7325):363-368.

Empain A. Estimation de la pollution par les métaux lourds dans la Somme par l'analyse des bryophytes aquatiques. *Bull. Fr. Piscicul.* 1976;260:138-142.

Evans JJ. Rubber tire leachates in the aquatic environment. *Rev. environ. Contam. Toxicol.* 1997;151:67-115.

Eyles J, Taylor SM, Johnson N, Baxter J. Worrying about waste : living close to solid waste disposal facilities in southern Ontario. *Soc.Scie.Med.* 1993;37(6):805-812.

Farquhar GJ, Rovers FA. Gas production during refuse decomposition. *Water, Air, Soil Pollution* 1973;2:483-493.

Feldmann J, Hirner AV. Occurrence of volatile metal and metalloid species in landfill and sewage gases. *Intern.J.EnvIRON.Anal.Chem.* 1995;60(2-4):339-359.

Fender H, Wolf G. Cytogenetic investigations in employees from waste disposal sites. *Toxicol. Lett.* 1998;96-97:148-154.

Fernandez JA, Aboal JR, Carballeira A. Use of native and transplanted mosses as complementary techniques for biomonitoring mercury around an industrial facility. *Sci. Total. Environ.* 2000;256:151-161.

Fielder HMP, Poon-King CM, Palmer SR, Moss N, Coleman G. Assessment of impact on health of residents living near the Nant-y-Gwyddon landfill site: retrospective analysis. *Br. med. J.* 2000;320:19-22.

Florjin PJ, Van Beusichem ML. Uptake and distribution of cadmium in maize inbred lines. *Plant Soil* 1993;150:25-32.

Fouchécourt MO, Rivière JL. Activities of liver and lung cytochrom P450 dependant

monoxygenases and antioxidant enzymes in laboratory and wild Norway rats exposed to reference and contaminated soils. Arch. Environ. Contam. Toxicol. 1996;30:513-522.

Foulks E, McLellen T. Psychologic sequelae of chronic toxic waste exposure. South Med.J. 1992;85(2):122-126.

France Nature Environnement. Bilan et analyse de la mise en place des Commissions Locales d'Information et de Surveillance; Décembre 1999. Rapport final.

Freeman NCG, Wainman T, Lioy PJ, Stern AH, Shupack SI. The effect of remediation of chromium waste sites on chromium levels in urine of children living in the surrounding neighbourhood. J. Air Waste Manage. Assoc. 1995;45(8):604-614.

Galsomiès L, Savanne D, Letrouit MA, Ayrault S, Charre B. Retombées atmosphériques des métaux en France : estimation par dosage dans les mousses. Campagne 1996. Paris; 1999. Ademe Editions.

Garrec JP, Van Haluwyn C. Biosurveillance végétale de la qualité de l'air. Editions TEC&DOC, Paris; 2002.

Geiger G, Federer P, Sticher H. Reclamation of heavy metal-contaminated soil : field studies and germination experiments. J. environ. Qual. 1993;22(1):201-207.

Gelberg KH. Health study of New York City Department of Sanitation landfill employees. J.Occup.Environ.Med. 1997;39(11):1103-1110.

Georgopoulos PG, Lioy PJ. Exposure measurement needs for hazardous waste sites: two case studies. Toxicol. ind. Health 1996;12:651-665.

GERAL. Unité de stockage d'ordures ménagères en zone rurale; 1998. Rapport Ademe.

GERAL. Environnement et santé : représentations des risques sanitaires liés aux déchets et à leurs modes de traitement: convention Ademe 970 8001; 1999.

Geschwind SA, Stolwijk JAJ, Bracken M, Fitzgerald E, Buckler G, Natarelli G et al. Risk of congenital malformations associated with proximity to hazardous waste sites. Am. J. Epidemiol. 1992;135(11):1197-1207.

Gochfeld M. Medical surveillance of hazardous waste workers: principles and problems. Occup. Med. 1990;5(1):1-7.

Gochfeld M, Udasin I, Favata E, Buckler G, Natarelli G, Burger J. Temporal changes in blood lead levels of hazardous waste workers in New Jersey, 1984-1987. Environ. Monit. Assess. 1993;25:99-107.

Goldberg MS, Goulet L, Riberdy H. Incidence of cancer among persons living near a municipal solid waste landfill site in Montreal, Quebec. Arch. environ. Health 1995b;50:416-424.

Goldberg MS, Goulet L, Riberdy H, Bonvalot Y. Low birth weight and preterm births among infants born to women living near a municipal solid waste landfill site in Montreal, Quebec. Environ. Res. 1995a;69:37-50.

Goldberg MS, Siemiatycki J, Dewar R, Desy M, Riberdy H. Risks of developing cancer relative to living near a municipal solid waste landfill site in Montreal, Quebec, Canada. *Arch. environ. Health* 1999;54(4):291-296.

Gombert S, Asta J. Etude du suivi de la pollution chlorée émise par une usine d'incinération d'ordures ménagères par l'utilisation de bioindicateurs végétaux : lichens et sphaignes. *Ecologie* 1997;28(4):365-372.

Gonsebatt ME, Salazar AM, Montero R, Diaz-Barriga F, Yanez L, Gomez H et al. Genotoxic monitoring of workers at a hazardous waste disposal site in Mexico. *Environ. Health Perspect.* 1995;103(1):111-113.

Gray PCR. Waste incineration: controversy and risk communication. *European review of Applied psychology* 1995;45(1):29-34.

Griffith J, McGeehin MA, Phifer BL, Ashley DL. Cancer mortality in U.S. countries with hazardous waste sites and ground water pollution. *Arch. environ. Health* 1989;44:69-74.

Grubinger VP, Gutenmann WH, Doss GJ, Rutzkz M, Lisk DJ. Chromium in Swiss chard grown on soil amended with tannery meal fertilizer. *Chemosphere* 1994;28(4):717-720.

Guidotti TL, Yoshida K, Clough V. Personal exposure to pesticide among workers engaged in pesticide container recycling operations. *Am. ind. Hyg. Assoc. J.* 1994;55(12):1154-1163.

Guillette IJ, Crain DA, Gross TS. Developmental abnormalities of the gonad and abnormal sex hormone concentrations in juvenile alligators from contaminated and control lakes in Florida. *Environ. Health Perspect.* 1994;102:680-688.

Hall HI, Price-Green PA, Dhara VR, Kaye WE. Health effects related to releases of hazardous substances on the superfund priority list. *Chemosphere* 1995;31(1):2455-2461.

Hamar GB, McGeehin MA, Phifer BL, Ashley DL. Volatile organic compound testing of a population living near a hazardous waste site. *J. Expo. Anal. environ. Epidemiol.* 1996;6(2):247-255.

Hara K, Hanaoka T, Yamano Y, Itani T. Urinary 1-hydroxypyrene levels of garbage collectors with low-level exposure to polycyclic aromatic hydrocarbons. *Sci. total Environ.* 1997;199(1-2):159-164.

Harding AK, Greer ML. The health impact of hazardous waste sites on minority communities: implications for public health and environmental health professionals. *J. Environ. Health* 1993;55(7):6-10.

Harnly M, Stephens R, McLaughlin C, Marcotte J, Petreas M, Goldman L. Polychlorinated dibenzo-p-dioxin and dibenzofuran contamination at metal recovery facilities, open burn sites, and a railroad car incineration facility. *Environ. Sci. Technol.* 1995;29(3):677-684.

Harrad SJ, Sewart P, Alcock R, Boumphrey R, Burnett, Duarte-Davidson R et al. Polychlorinated biphenyls (PCBs) in the British environment : sinks, sources and temporal trends. *Environ. Pollut.* 1994;85:131-146.

Hartmann A, Fender H, Speit G. Comparative biomonitoring study of workers at a waste disposal site using cytogenetic tests and the comet (single-cell gel) assay. *Environ. Mol.*

Mutagen. 1998;32:17-24.

Hawksworth DL, Rose F. Qualitative scale for estimating sulphur dioxide air pollution in England and Wales using epiphytic lichens. *Nature* 1970;227:145-148.

Heederik D, Douwes J. TOwards an occupational exposure limit for endotoxins? *Ann. Agric. Environ. Med.* 1997;4:17-19.

Hertzman C, Hayes M, Singer J, Highland J. Upper Ottawa street landfill site health study. *Environ. Health Perspect.* 1987;75:173-195.

Herzig R, Liebendörfer L, Urech M. Flechten als bioindikatoren der Luftverschmutzung in der Schweiz : Methoden-Evaluation und Eichung mit wichtigen Luftschadstoffen. *VDI Berichte* 1987;609:619-639.

Horvath B, Gruiz K. Impact of metalliferous ore mining activity on the environment in Gyongyosorosi, Hungary. *Sci. Total. Environ.* 1996;184:215-227.

Hoskin AF, Leigh JP, Planek TW. Estimated risk of occupational facilities associated with hazardous waste site remediation. *Risk Anal.* 1994;14(6):1011-1017.

Hours M, Anzivino L, Herrscher MP, Perrodin Y, Sarrazin B, Maitre A et al. Mise au point d'une méthodologie de surveillance des polluants atmosphériques dans les centres de stockage des ordures ménagères : application à deux sites. Réseau Santé Déchets. Financement RE.CO.R.D., SITA, CREED, Ademe; déc. 2000. 2000a. 243p+annexes.

Hours M, Anzivino L, Asta J, Berny P, Billeret M, Maitre A et al. Etude des polluants atmosphériques émis dans deux centres de stockage des ordures ménagères: Réseau Santé Déchets. Financement RE.CO.R.D., SITA, CREED, Ademe; nov. 2000. Rapport final. 2000b.

Hours M, Anzivino L, Asta J, Billeret M, Berny P, Keck G et al. Etude des polluants atmosphériques émis dans deux centres de stockage des ordures ménagères : caractérisation et mesure des niveaux d'exposition : Mise au point d'outils de suivi en vue de l'évaluation des risques sanitaires: Réseau Santé Déchets; 2000c. Rapport final. Tome 1 : 376p. Tome 2 (annexes) : 114p.

Hours M, Perrodin Y, Asta J, Berny P, Parat S, Stoklov M et al. Etude des polluants atmosphériques émis dans deux centres de stockage des ordures ménagères : caractérisation et mesure des niveaux d'exposition ; mise au point d'outils de suivi en vue de l'évaluation des risques sanitaires. *Déchets, sciences et techniques* 2001;24:38-43.

Huber MS, Gerba CP, al. Study of Persistence of Enteric Viruses in Landfilled disposable diapers. *Environ. Sci. Technol.* 1994;28:1767-1772.

Hyslop BT, Davies MS, Artur W, Gazey NJ, Holroyd S. Effects of colliery waste on littoral communities in North-East England. *Environ. Pollut.* 1997;96(3):383-400.

Ineris. Amélioration de la gestion des décharges d'ordures ménagères et assimilées sous l'angle de leur impact environnemental ; 1996. Rapport final. Auteur Bour O. 23p.+annexes.

Ineris. Analyses des rejets atmosphériques de torchères de biogaz: Ministère de l'Environnement, Ademe, SNAD; Juillet 1999. Rapport Final. Auteur: Van Elsuve D.

INRS. Cahier de notes documentaires n°156. 1994.

Jacobson JS, Hill AC. Recognition of air pollution injury to vegetation : a pictorial atlas. Pittsburgh, Pennsylvania: Air Pollution Control Association; 1970. Informative Report 1, TR-7 Agricultural Committee.

Jakubowski M, Trzcinka-Ochocka M, Razniewska G, Matczak W. Biological monitoring of occupational exposure to arsenic by determining urinary content of inorganic arsenic and its methylated metabolites. *Int. Arch. occup. environ. Health* 1998;71 Suppl:S29-32.

James WH. Offspring sex ratio as a potential monitor of reproductive disorders in communities near hazardous chemical sites. *Reproduct. Toxicol.* 1997;11(6):893-895.

Jarup L, Briggs D, de Hoogh C, Morris S, Hurt C, Lewin A et al. Cancer risks in populations living near landfill sites in Great Britain. *Br. J. Cancer* 2002;86(11):1732-1736.

Jarup L, Persson B, Elinder CG. Blood cadmium as an indicator of dose in a long-term follow-up of workers previously exposed to cadmium. *Scand. J. Work environ. Health* 1997;23(1):31-36.

Jenner HA, Janssen-Mommen JPM. Duckweed *Lemna minor* as a tool for testing toxicity of coal residues and polluted sediments. *Arch. Environ. Contam. Toxicol.* 1993;25:3-11.

Johnson BL. A review of the effects of hazardous waste on reproductive health. *Am. J. Obstetrics Gynecol.* 1999;181(1):S12-S16.

Johnson BL, Coulberson SL. Environmental epidemiologic issues on minority health. *AEP* 1993;3(2):223-232.

Johnson BL, Grandjean P, Amler RW. Neurobehavioral testing and hazardous chemical sites. *Neurotoxicol. Teratol.* 1994;16(5):485-487.

Johnson MS, Leah RT, Connor L, Rae C, Saunders S. Polychlorinated biphenyls in small mammals from contaminated landfill sites. *Environ. Pollut.* 1996;92:195-191.

Junyapoon S, Bartle KD, Ross AB, Cooke M. Analysis of malodorous sulfur gases and volatile organometalloïd compounds in landfill gas emissions using capillary gas chromatography with programmed temperature vaporization injection and atomic emission detection. *Intern.J.Envirion.Anal.Chem.* 2002;82(2):47-59.

Kharrazi M, von Behren J, Smith M, Lomas T. A community-based study of adverse pregnancy outcomes near a large hazardous waste landfill in California. *Toxicol. ind. Health* 1997;13(2-3):299-310.

Kim S, Stair E, Lochmiller RL, Rafferty DP, Schroder JL, Basta NT et al. Widespread risks of dental fluorosis in cotton rats (*Sigmodon hispidus*) residing on petrochemical waste sites. *J. Toxicol. environ. Health* 2001;62(2):107-125.

Kirivanta H, Tuomainen A, Reiman M, Laitinen S, Nevalainen A, Liesivuori J. Exposure to airborne microorganisms and volatile organic compounds in different types of waste handling. *Ann. Agric. Environ. Med.* 1999;5:39-44.

Kjeldsen P, Barlaz MA, Rooker AP, Baun A, Ledin A, Christensen TH. Present and long-term

composition of MSW landfill leachate: a review. *Crit.Rev.Environ.Sci.Technol.* 2002;32(4):297-336.

Kjeldsen P, Christophersen M. Composition of leachate from old landfills in Denmark. *Waste Manag.Res.* 2001;19(3):249-256.

Klemans W, Vleminckx C, Schriewer L, Joris I. Cytogenetic biomonitoring of a population of children allegedly exposed to environmental pollutants. Phase 2: results of three-year longitudinal study. *Mutat. Res.* 1995;342:147-156.

Kocan A, Bencko V, Sixl W. Polychlorinated dibenzo-p-dioxins (PCDDs) and dibenzofurans (PCDFs) in the hair of people living on municipal refuse dumping sites in Cairo (Egypt). *Toxicol. environ. Chem.* 1992;36:33-37.

Kruckeberg AL, Wu L. Copper tolerance and copper accumulation of herbaceous plants colonizing inactive California copper mines. *Ecotoxicol. environ. Saf.* 1992;23:307-319.

Krug MN, Ham RK. Analysis of long-term leachate characteristics. In: *Proceedings Sardinia 1997, 6th International Landfill Symposium*; S. Margherita di Pula, Cagliari, Italy; 1997. p. 117-132.

Lagadic L, Caquet T, Amiard JC. Biomarqueurs en écotoxicologie : principes et définitions. In: Masson, editor. *Lagadic L., Caquet T., Amiard J.C. & Ramade F. Biomarqueurs en écotoxicologie. Aspects fondamentaux.* Paris; 1997. p. 1-9.

Lakhanisky T, Bazzoni D, Jadot P, Joris I, Laurent C, Ottogali M et al. Cytogenetic monitoring of a village population potentially exposed to a low level of environmental pollutants. Phase 1: SCE analysis. *Mutat. Res.* 1993;319:317-323.

Lange JH, Thomulka KW. Disposal of hazardous waste: cost and liability issues. *Fresenius environ. Bull.* 1995;4:303-308.

Lascoumes P. "La précaution comme anticipation des risques résiduels et hybridation de la responsabilité". In: *Ecole des Mines de Paris, Centre de sociologie de l'innovation. Représenter, hybrider, coordonner, Actes des journées des 9 et 10 mai 1996*; 1996; Paris.

Laurent C, Lakhanisky T, Jadot P, Joris I, Ottogali M, Planard C et al. Increased sister chromatid exchange frequencies observed in a cohort of inhabitants of a village located at the boundary of an industrial dumping ground: phase 1. *Cancer Epidemiol. Biomarkers Prev.* 1993;2:355-362.

Lavoie J, Guertin S. Evaluation of health and safety risks in municipal solid waste recycling plants. *J. Air Waste Manage. Assoc.* 2001;51:352-360.

Leblanc F, De Sloover J. Relation between industrialization and the distribution and growth of epiphytic lichens and mosses in Montreal. *Canad. Jour. Bot.* 1970;48:1485-1496.

Ledrans M, Boudot J. Surveillance du saturnisme infantile en France. *Bilan des activités de dépistage*: Institut de Veille Sanitaire; 1998.

Leigh JP, Hoskin AF. Hazards for nearby residents and cleanup workers of waste sites. *J. occup. environ. Med.* 1999;41(5):331-348.

Levine SP, Turpin SD, Gochfeld M. Protecting personnel at hazardous waste sites: current issues. *Appl. occup. environ. Hyg.* 1991;6(12):1007-1014.

Lewis-Michl EL, Kallenbach LR, Gearyns, Melius JM, Ju CL, Orr MF et al. Investigation of cancer incidence and residence near 38 landfills with soil gas migration conditions, New York State, 1980 - 1989. Atlanta, Georgia: Agency for Toxic Substances and Disease Registry; 1998. Report for grant #H75/ATH298290-01 to the Department of Health and Human Sciences.

Lhuillier D, Cochin Y. *Des déchets et des hommes*: Editions Desclée de Brower, collection "sociologie clinique; 1999. 185p.

Linder G, Bollman M, Gillet C, King R, Nwosu J, Ott S et al. A framework for field and laboratory studies for ecological risk assessment in wetland and terrestrial habits : two case studies. In: Hughes J.S., Biddinger, G.R. & Mones E. (eds), editor. *Environmental toxicology and risk assessment*. ATSM Symposium; April 26-29 1993; Atlanta; 1993. p. 61-78.

Lioy PJ. Measurement methods for human exposure analysis. *Environ. Health Perspect. Suppl.* 1995;103(3):35-43.

Lipscomb J, Goldmann LR, Satin K, Smith DF, Vance WA, Neutra RR. A follow-up study of the community near the Mc Coll waste disposal site. *Environ. Health Perspect.* 1991;94:15-24.

Llobet JM, Schuhmacher M, Domingo JL. Spatial distribution and temporal variation of metals in the vicinity of a municipal solid waste incinerator after a modernization of the flue gas cleaning systems of the facility. *Sci. total Environ.* 2002;284(1-3):205-214.

Lober DJ. Beyond self-interest: a model of public attitudes towards waste facility siting. *J. Environ. Plan. Manage.* 1993;36(3):345-363.

Lober DJ. Why protest? Public behavior and attitudinal response to siting a waste disposal facility. *Policy Stud. J.* 1995;23:499-518.

Lober DJ, Green DP. Nimby or niaby: a logit model of opposition to solid-waste-disposal facility siting. *J. environ. manage.* 1994;40:33-50.

Locett Doust J, Schmidt M, Locett Doust L. Biological assessment of aquatic pollution : a review, with emphasis on plants as biomonitors. *Biol. Rev.* 1994;69:147-186.

Logan TJ, Goins LE, Lindsay BJ. Field assessment of trace element uptake by six vegetables from N-Viro soil. *Water Environ. Res.* 1997;69(1):28-33.

Loizidou M, Kapetanios EG. Study on the gaseous emissions from a landfill. *Sci. total Environ.* 1992;127:201-210.

Luotamo M, Patterson DG, Needham LL, Aitio A. Concentrations of PCB congeners in sera from workers with past and present exposure. *Chemosphere* 1993;27(1-3):171-177.

Luskin J, Somers C, Wooding J, Levenstein C. Teaching health and safety: problems and possibilities for learner-centered training. *Am. J. ind. Med.* 1992;22:665-676.

Makridis C, Pateras D, Amberer A. Thallium pollution risk to food chain from cement plant.

Fresenius environ. Bull. 1996;5(11-12):643-648.

Mallin K. Investigation of a bladder cancer cluster in northwestern Illinois. Am. J. Epidemiol. 1990;132:S96-S106.

Margai FL. Evaluating the potential for environmental quality improvement in a community distressed by manmade hazards. J.envIRON.MANAGE. 1995;44(2):181-190.

Marshall EG, Geensburg LJ, Deres DA, Geary NS, Cayo MR. Maternal residential exposure to hazardous wastes and risk of central nervous system and musculoskeletal birth defects. Arch. environ. Health 1997;52(6):416-425.

Marth E, Sixl W, Bencko V, Medwed M, Lapajne S, Voncina E et al. People on the garbage dumps of Cairo: a toxicological in vivo model ? Cent. Eur. J. public Health 1995;3:154-157.

McCarron P, Harvey I, Brogan R, Peters TJ. Self reported health of people in an area contaminated by chromium waste: interview study. Br. med. J. 2000;320(7226):11-15.

McCullough JE, Henderson AK, Kaufman JD. Occupational burns in Washington State, 1989-1993. J. occup. environ. Med. 1998;40(12):1083-1089.

McDiarmid MA, Freeman CS, Grossman EA, Martonik J. Biological monitoring results for cadmium exposed workers. Am. ind. Hyg. Assoc. J. 1996;57(11):1019-1023.

McGrath SP, Chang AC, Page AL, Witter E. Land application of sewage sludge: scientific perspectives of heavy metal loading limits in Europe and the United States. Environ. Rev. 1994;2:108-118.

McQuiston TH, Coleman P, Wallerstein NB, Marcus AC, Morawetz JS, Ortlieb DW. Hazardous waste worker education: long-term effects. J.Occup. Med. 1994;36(12):1310-1323.

Melius JM. Medical surveillance for hazardous waste workers. J.Occup. Med. 1986;28(8):679-683.

Melius JM. OSHA standard for medical surveillance of hazardous waste workers. Occup. Med. 1990;5(1):143-150.

Meneses M, Llobet JM, Granero S, Schuhmacher M, Domingo JL. Monitoring metals in the vicinity of a municipal waste incinerator: temporal variation in soils and vegetation. Sci. total Environ. 1999;226:157-164.

Millot N. Les lixiviats de décharges contrôlées : caractérisation analytique. Etude des filières de traitement. Thèse INSA de Lyon (LCPAE); 1986. 188p.

Mozzon D, Brown DA, Smith JW. Occupational exposure to airborne dust, respirable dust and metals arising from refuse handling, burning and landfilling. Am. J. Epidemiol. 1987;48(2):111-116.

Najem LR, Strunck T, Feuerman M. Health effects of a superfund hazardous chemical waste disposal site. Am. J. prev. Med. 1994;10(3):151-155.

Nedellec V, Mosqueron L. Recensement des agents émis lors des déversements d'ordures

ménagères en situation professionnelle et identification des dangers par inhalation. *Environnement, Risques et Santé* 2002;1(3):164-177.

Neutra RR, Lipscomb J, Satin K, Shusterman D. Hypotheses to explain the higher symptom of the community near the McColl waste disposal site. *Environ. Health Perspect.* 1991;94:31-38.

Nimis PL. Linee-guida per la bioindicazione degli effetti dell'inquinamento tramite la biodiversità dei licheni epifiti. In: Piccini C. & Salvati S., editor. Proc. Workshop " Biomonitoraggio della qualità dell'aria sull territorio nazionale "; 26-27 November 1998; Roma; 1999a. p. 267-277.

Nimis PL. The future of the bio-indication with lichens. *International Lichenological Newsletter* 1999b;32.

Nouwen J, Cornelis C, De Fre R, Wevers M, Viane P, Mensink C et al. Health risk assessment of dioxin emissions from municipal waste incinerators: the Neerland quarter (Wilrijk, Belgium). *Chemosphere* 2001;43:909-923.

Nylander W. Les lichens du Jardin du Luxembourg. *Bull. Soc. Bot.* 1866;13:364-372.

Nylander W. Les lichens des environs de Paris. Ed. Schmidt. Paris; 1896. 182p.

Okrent D, Xing L. Future risk from a hypothesized RCRA site disposing of carcinogenic metals should a loss of societal memory occur. *J. hazard. Mater* 1993;38:363-384.

Osterman-Golkar SM, Macneela JP, Turner MJ, Walker VE, Swenberg JA, Sumner SJ et al. Monitoring exposure to acrylonitrile using adducts with N-terminal valine in haemoglobin. *Carcinogenesis* 1994;15(12):2701-2707.

Ostry AS, Hertzman C, Teschke K. Community risk perception : a case study in a rural community hosting a waste site used by a large municipality. *Canadian J. of public health* 1993;84(6):415-418.

Ostry AS, Hertzman C, Teschke K. Community risk perception and waste management : a comparison of three communities. *Arch. Environ. Health* 1995;50(2):95-102.

Pakeman RJ, Hankard PK, Osborn D. Plants as biomonitors of atmospheric pollution : their potential for use in pollution regulation. *Rev. environ. Contam. Toxicol.* 1998;157:1-23.

Parker T, Dottridge J, Kelly S. Investigation of the composition and emissions of trace components in landfill gas: Environmental Agency, Royaume Uni; 2002. R&D Technical Report P1-438/TR.

Paustenbach D, Burke ML, Shum M, Kalmes R. Airborne concentrations of ethyl and methyl cyanoacrylate in the workplace. *AIHAJ: J. Sci. occup. environ. I Health Saf.* 2001;62(1):70-79.

Pedersen BA, Higgins GM. Evaluation of chemical exposures in the hazardous waste industry. *J. Air Waste Manage. Assoc.* 1995;45(2):89-94.

Perdrix A, Madon N, al. Risques biologiques autres qu'infectieux. In: Encyclopédie médico-chirurgicale, Toxicologie-Pathologie professionnelle 2; (16-080-B-10) ;1997. p. 1-6.

Peretti-Wattel P. Sociologie du risque; A. Colin, Paris 2000.

Petts J. Incineration risk perception and public concern : experience in the U.K. improving risk communication. Waste management research 1992;10:169-182.

Petts J. Effective waste management: understanding and dealing with public concerns. Waste management research 1994;12:207-222.

Piechaczyk X. Instruire ou construire l'intérêt général? Radiographie d'une population de commissaires enquêteurs. Ecologie et politique 1997; n° 21.

Pihan F, De Vaufleury A. The snail as a target organism for the evaluation of industrial waste dump contamination and the efficiency of its remediation. Ecotoxicol. environ. Saf. 2000;B46:137-147.

Pinel C, Parat S et al. Mycotoxines. In: *Encyclopédie Médico-chirurgicale. Toxicologie-Pathologie professionnelle*; (16-070-A-10) ;1998. p. 1-5.

Pohland FG, Harper SR. Critical review and summary of leachate and gas production from landfills.: Georgia Institute of Technology (Atlanta, USA); 1985. Project n°E-20-G01, EPA CR 809 997.

Poulleau J. Caractérisation des biogaz. Bibliographie. Mesures sur site. Ineris; 2002. Rapport final.

Poulsen OM, Breum NO et al. Sorting and recycling of domestic waste. Review of occupational health problems and their possible causes. The Science of the Total Environment 1995;168:33-56.

Pronost R, Matejka G. Les lixiviats de décharges d'ordures ménagères. Production, caractérisation, traitement. Environnement & Technique 2000;196:25-29.

Propst TL, Lochmiller RL, Qualis CWJR, McBee K. In situ (mesocosm) assessment of immunotoxicity risks to small mammals inhabiting petrochemical waste sites. Chemosphere 1999;38(5):1049-1067.

Rahkonen P, Ettala M. Airborne microbes and endotoxins in the work environment of two sanitary landfills in Finland. Aerosol Sciences and Technology 1990;13:505-513.

Rahkonen P, Malmen Y, Ettala M. Work safety in collection and transport of hazardous wastes. Waste Manage. Res. 1992a;10:217-226.

Ramade F. Précis d'Ecotoxicologie. Ed. Masson. Paris; 1992. 300p.

Ramade F. Dictionnaire encyclopédique de l'écologie et des sciences de l'environnement. Ediscience, Paris; 1993. 822p.

Ramade F. Dictionnaire encyclopédique des pollutions. Ediscience Paris; 2000. 690p.

Raman N, Sambandan K. Distribution of VAM Fungi in tannery effluent polluted soils of Tamil Nadu, India. Bull. Environ. Contam. Toxicol. 1998;60:142-150.

Reams MA, Templet PH. Political and environmental equity issues related to municipal waste

incineration siting. *J.Hazard.Mater.* 1996;47:313-323.

Rees JF. The fate of organic compounds in the landfill disposal of organic matter. *Journal of Chem. Tech. Biotechnol.* 1980;30:161-175.

Reif JS, Tsongas TA, Anger WK, Mitchell J, Metzger L, Keefe TJ et al. Two-stage evaluation of exposure to mercury and biomarkers of neurotoxicity at a hazardous waste site. *J. Toxicol. environ. Health* 1993;40:413-422.

Reinhart DR. A review of recent studies on the sources of hazardous compounds emitted from solid waste landfills : a U.S. experience. *Waste Manag.Res.* 1993;11(3):257-268.

Reinthal FF, Haas D, al. Comparative investigations of airborne culturable microorganisms in selected waste treatment facilities and in neighbouring residential areas. *Zentralbl Hyg Umweltmed* 1999;202(1):1-17.

Rettenberger G, Stegmann R. Landfill gas components. In: E&FN Spon, editor. *Landfilling of waste : biogas.* Londres; 1996.

Rico A. Chemo-defence system. *C.R. Acad. Sci. Paris. Sciences de la vie/Life Sciences* 2001;324:97-106.

Rivière JL. Evaluation du risque écologique des sols pollués. *Record Lavoisier Tec&Doc,* Paris; 1998. 230p.

Romine JD, Barth EF. Preliminary assessment of worker and ambient air exposures during soil remediation technology demonstration. *Am. ind. Hyg. Assoc. J.* 2002;63(1):76-80.

Roos PH, Van Afferden M, Strotkamp D, Tappe D, Pfeifer F, Hanstein WG. Liver microsomal levels of cytochrome P4501a1 as biomarker for exposure and bioavailability of soil-bound polycyclic aromatic hydrocarbons. *Arch. Environ. Contam. Toxicol.* 1996;30(1):107-113.

Rout GR, Samantaray S, Das P. Effects of chromium and nickel on germination and growth in tolerant and non-tolerant populations of *Echinochloa colona* (L.) Link. *Chemosphere* 2000;40:855-859.

Rühling A, Tyler G. An ecological approach to the lead problem. *Bot. Notiser* 1968;121:321-342.

Ruokojärvi P, Ettala M, Rahkonen P, Tarhanen J, Ruuskanen J. Polychlorinated dibenzo-p-dioxins and -furans (PCDDs and PCDFs) in municipal waste landfill fires. *Chemosphere* 1995;30(9):1697-1708.

Salkin IF. Review of health impacts from microbiological hazards in health-care wastes. *WHO Geneva* 2001:1-25.

Sauve S, Cook N, Hendershot WH, McBride MB. Linking plant tissue concentrations and soil copper pools in urban contaminated soils. *Environ. Pollut.* 1996;94(2):153-157.

Schrive SC, McBride RA. Physiological responses of red maple saplings to sub-irrigation with an untreated municipal landfill leachate. *Waste Manage. Res.* 1995;13(3):219-239.

Schuhmacher M, Granero S, Rivera J, Müller L, Llobet JM, Domingo JL. Atmospheric

deposition of PCDD/Fs near an old municipal solid waste incinerator: levels in soil and vegetation. *Chemosphere* 2000;40:593-600.

Schulte PA. Opportunities for the development and use of biomarkers. *Toxicol. Lett.*;77(1-3):25-29.

Schweisfurth H, Schöttes C. Acute intoxication of a hydrazine-like gas by 19 workers in a garbage dump. Akute Intoxikation von hydrazinähnlichem gas bei 19 Arbeitern einer Mülldeponieanlage. *Zent. bl. Hyg. Umweltmed.* 1993;195:46-54.

Scott P, MacDonald C. The significance and environmental impact of trace compounds present in gaseous emissions from U.K. landfill sites. *Tiré à part* 1990:10p.

SFSP (Société Française de Santé Publique). L'incinération des déchets et la santé publique : bilan des connaissances récentes et évaluation du risque; Collection santé et société n°7. 1999. 368p.

Shusterman D, Lipscomb J, Neutra RR, Satin K. Symptom prevalence and odor-worry interaction near hazardous waste sites. *Environ. Health Perspect.* 1991;94:25-30.

Sillet AA, Royer S, Coque Y, Thomas O. Les lixiviats de décharges d'ordures ménagères : genèse, composition et traitement. *Déchets Sciences & Techniques* 2001;22:7-11.

Smits JE, Wayland ME, Miller MJ, Liber K, Trudeau S. Reproductive, immune, and physiological end points in tree swallows on reclaimed oil sands mine sites. *Environ. Toxicol. Chem.* 2000;19:2951-2960.

Sosniak WA, Kaye WE, Gomez TM. Data linkage to explore the risk of low birthweight associated with maternal proximity to hazardous waste sites from the national priorities list. *Arch. environ. Health* 1994;49(4):251-255.

Stahl RG. Can mammalian and non-mammalian "sentinel species" data be used to evaluate the human health implications of environmental contaminants? *Hum. ecol. Risk Assess.* 1997;3(3):329-335.

Steer PJ, Tashiro CHM, Mcillveen WD, Clement RE. PCDD and PCDF in air, soil, vegetation and oily runoff from a tire fire. *Water Air Soil Pollut.* 1995;82(3-4):659-674.

Stern AH, Fagliano JA, Savrin JE, Freeman NCG, Liroy PJ. The association of chromium in household dust with urinary chromium in residences adjacent to chromate production waste sites. *Environ. Health Perspect.* 1998;106(12):833-839.

Stern AH, Freeman NCG, Pleban P, Boesch RR, Wainman T, Howell T et al. Residential exposure to chromium waste - urine biological monitoring in conjunction with environmental exposure monitoring. *Environ. Res.* 1992;58:147-162.

Suurnakki T, Ilmarinen J, Wagar G, Jarvinen E, Landau K. Municipal employees' cardiovascular diseases and occupational stress factors in Finland. *Int. Arch. occup. environ. Health* 1987;59(2):107-114.

Tarkowski S, Rolecki R. Hazardous wastes and public health. Potential health effects of exposure at levels encountered around the landfills. *Eur. Epimarcker* 2001;5(2):1-7.

Thimonier A, Dupouey JL, Bost F, Bexker M. Simultaneous eutrophication and acidification of a forest ecosystem in North-East France. *New Phytol.* 1994;126:533-539.

Touze-Foltz N. Détection, localisation et contrôle des fuites dedans les géomembranes. *Environnement & Technique* 2003;223:21-25.

Udasin IG, Buckler G, Gochfeld M. Quality assurance audits of medical surveillance programs for hazardous waste workers. *J.Occup. Med.* 1991;33(11):1170-1174.

Udger DG, Wandersman A, Hallman W. Living near a hazardous waste facility: coping with individual and family distress. *Am.J.Orthopsychiatr.* 1991;62(1):55-70.

Ulfig K, Plaza G, Szytyler A, Bronder J, Terkowski M, Guarro J. General assessment of the influence of a municipal landfill site and environmental factors on the occurrence of keratolytic fungi in soil. *Roczn. Panstwowego Zakl. Hig.* 2000;51(2):167-181.

Van Tongeren M, Van Amelsvoort L et al. Exposure to organic dusts, endotoxins, and microorganisms in the municipal waste industry. *Int. J. of Environmental Health* 1997b;3(Jan-Mar):30-36.

Van Tongeren M, Van Amelsvoort L, Heederik D. Exposure to organic dusts, endotoxins, and microorganisms in the municipal waste industry. *Int. J. occup. Environ. Health* 1997a;3(1):30-36.

Vasseur L, Fortin MJ, Cyr J. Clover and cress as indicator species of impacts from limed sewage sludge and landfill wastewater land application. *Sci. Total. Environ.* 1998;217(3):231-239.

VDI. Measurement of immission effects. Measurement and evaluation of phytotoxic effects of ambient air pollutants (immissions) with lichens. Mapping of lichens for assessment of the air quality. *Richtlinien*; 1995; 3799.

Verneaux J, Tuffery G. Une méthode zoologique pratique de détermination de la qualité biologique des eaux courantes. *Indices biotiques. Annal. Scient. Univ. Besançon, Zoologie* 1967;3:79-90.

Vrijheid M. Health effects of residence near hazardous waste landfill sites: a review of epidemiologic literature. *Environ. Health Perspect.* 2000;108(suppl1):101-112.

Vrijheid M, Dolk H, Armstrong B, Abramsky L, Bianchi F, Fazarinc I et al. Chromosomal congenital anomalies and residence near hazardous waste landfill sites. *Lancet* 2002;359(9303):320-322.

Wakefield S, Elliott SJ. Environmental risk perception and well-being : effects of the landfill siting process in two southern Ontario communities. *Soc.Scie.Med.* 2000;50:1139-1154.

Wang W, Freemark K. The use of plants for environment monitoring and assessment. *Ecotoxicol. environ. Saf.* 1995;30(3):289-301.

White RF, Feldman RG, Eviator II, Jabre JF, Niles CA. Hazardous waste and neurobehavioral effects: a developmental perspective. *Environ. Res.* 1997;73:113-124.

Whiteman DC, Dunne MP, Burnette PC. Psychological and social correlates of attrition in a

longitudinal study of hazardous waste exposure. Arch.Environ.Health 1995;50(4):281-286.

Wickliffe JK, Pitts R, Iverson S, Bickham JW. Cell cycle disruption in wild rodent populations as an endpoint in detecting exposure and effect. Bull. Environ. Contam. Toxicol. 2000;64:448-454.

Williams A, Jalaludin B. Cancer incidence and mortality around a hazardous waste depot. Aust. N. Z. J. Public Health 1998;22(3):324-346.

Winner WE. Responses of bryophytes to air pollution. In: Cramer, editor. Nash III TH. & Wirth V. Lichens, Bryophytes and air quality. Berlin; 1988. p. 141-173.

Xiong ZT, Hu HX, Wang YX, Fu GH, Tan ZQ, Yan GA. Comparative analyses of soil contaminant levels and plant species diversity at developing and disused oil well sites in Qianjiang oilfield, China. Bull. Environ. Contam. Toxicol. 1997;60(4):667-672.

Zaman MS, Zereen F. Growth responses of radish plants to soil cadmium and lead contamination. Bull. Environ. Contam. Toxicol. 1998;61:44-50.

Zhu YG, Smolders E. Plant uptake of radiocaesium : a review of mechanisms, regulation and application. J. exp. Bot. 2000;51(351):1635-1645.

Zirschky J, Gentry B, Marcus P. Superfund and contamination of workers' homes. Am. ind. Hyg. Assoc. J. 1987;48(11):A718, A720, A722-724.

Zmirou D, Dab W. Montchanin: retour d'expérience. In: Editorial pour Info Santé Déchets n°22. Lyon: Réseau Santé Déchets; 1998.

Zmirou D, Deloraine A, Dab W, Isnard H. Incertitude, expertise et décision: les leçons de la décharge de Montchanin. Rev. Epidém. et Santé Publ. 1995;43:584-593.

Zmirou D, Deloraine A, Saviuc P, Tillier C, Boucharlat A, Maury N. Short-term health effects of an industrial toxic waste landfill: a retrospective follow-up study in Montchanin, France. Arch. environ. Health 1994;49(4):228-238.

Zmirou D, Perrodin Y. Approche méthodologique de l'évaluation des risques pour la santé des déchets stabilisés. Application à un scénario. Déchets Sci. Tech. 1999;16:10-13.

CHAPITRE 3 EVALUATION DES RISQUES SANITAIRES CHRONIQUES

Agences de l'Eau, Ministère de l'Environnement. Guide pratique pour le contrôle et l'entretien des captages d'eau souterraine; 1994. 40p.

Agences de l'Eau, Ministère de l'Environnement. Etude inter-Agences de l'eau : Evaluation des flux polluants liquides générés par les décharges de déchets ménagers; 1997.

Assmuth T. Concentrations of toxic chemicals in waste deposit runoff. *Aqua Fennica* 1992;21(2):183-194.

Assmuth T. Comparative risk analysis of waste site toxicants by indices based on concentration distributions, fluxes, environmental fate and critical effects. *J Hazardous Materials* 1996;48:121-135.

ATSDR (Agency for Toxic Substances and Disease Registry). 1999 CERCLA priority list of hazardous substances that will be the subject of toxicological profiles & support document: Division of toxicology, US Department of Health and human services; octobre 1999.

ATSDR (Agency for Toxic Substances and Disease Registry). 2001 CERCLA priority list. 2002.

Beaudeau P, Zeghnoun A, Ledrans M, Volatier JL. Consommation d'eau du robinet pour la boisson en France métropolitaine : valeurs de référence pour l'épidémiologie et l'évaluation quantitative des risques sanitaires sur la base de l'enquête INCA1. *Environnement, Risques et Santé* 2003;2(3):147-158.

Belevi H, Baccini P. Long-term behavior of municipal solid waste landfills. *Waste management research* 1989;7(43).

Bendz D, Flyhammar P. Channel flow and its effects on long-term leaching of heavy metals in MSW landfills. In: *Seventh International Waste Management and Landfill Symposium*; 1999;2:43-50.

Bertrand M. Consommation et lieux d'achat des produits alimentaires en 1991. *Insee Résultats* 1993;262-263(54-55).

Bonaparte R, Gross BA. Field behavior of double-liner systems in waste containment facilities. *American Society of Civil Engineers; Geotechnical special publication 26* ;1990.

Bonnard R. Evaluation de l'impact sur la santé des rejets atmosphériques des tranches charbon d'une grande installation de combustion. Partie 2 : Exposition par voies indirectes: Ineris; 2003. DRC-03-45956/ERSA-n°92-RBn-/GT-GICversion4.

California Environmental Protection Agency. CalTOX A multimedia total exposure model for hazardous waste sites. Sacramento, California.; 2002.

Chiriack R. Etude des émissions des COV par les Centres d'Enfouissement Techniques d'OM. Evaluation de l'impact environnemental: Thèse Ademe UCB Lyon 1; 2002.

Christensen TH, Cossu R, Stegmann R. *Landfilling of Waste: Biogas*. E&FN SPON. London; 1996.

Coppalle A, Beaudeau P, Delmas V. Le transfert extérieur/intérieur des polluants atmosphériques: approches statistiques et physiques utiles pour le calcul de l'exposition des populations. *Pollution atmosphérique* 2002;175:407-419.

CREDOC, AFSSA, Ministère de l'Agriculture et de la Pêche. Enquête INCA individuelle et

nationale sur les consommations alimentaires: Editions TEC & DOC; 2000.

Dor F. Santé-Déchets: Certitudes et Incertitudes. Environnement et Technique 1998;176:27-29.

Dor F, Bonvallet N. Insecticides Organochlorés aux Antilles : Identification des dangers et Valeurs Toxicologiques de Référence (VTR) - Etat des connaissances et pertinence de l'utilisation dans la démarche d'EQRS. Institut de Veille Sanitaire Avril 2003, 56 p.

Duffa C, Delery L. Concentrations en substances chimiques dans l'environnement terrestre du Nord-Cotentin. Revue des données sur les niveaux moyens rencontrés dans l'environnement: Ineris; 2002. Rapport DPRE/SERNAT/ 02-10.

ECETOC. Exposure Factors Sourcebook for European Populations (with Focus on UK Data). Brussels; juin 2001. n° 79.

Feuillade G, Lagier T, Matejka G. Les métaux lourds dans les décharges d'ordures ménagères : synthèse des connaissances. Partie II : synthèse des connaissances. Tribune de l'eau 2001(611-613):25-36.

Gagnon F, Avard D, Ayotte P, Gingras B, Grondin J, Harmsen E et al. Groupes à risques particuliers. In: TEC&DOC, editor. Gérin M. et al. "Environnement et Santé Publique. Fondements et pratiques." Canada; 2003. 1023p.

Gibbons RD, Dolan DG, May H, O'Leary K, O'Hara R. Statistical comparison of leachate from hazardous, codisposal, and municipal solid waste landfills. Ground Water Monit.Rev. 1999;19(72).

Groupe Radioécologie Nord-Cotentin. Evaluation des risques pour la santé humaine associés aux rejets chimiques des installations nucléaires du Nord-Cotentin: IRSN; 2002.

Guyonnet D. MISP_v1. Un modèle analytique pour estimer l'impact de sources de pollution sur les eaux souterraines. Guide d'utilisation; 2001. Rapport BRGM RP-51039-FR.

Hjelmar O, Van der Sloot HA, Guyonnet D, Rietra RPJJ, Brun A, Hall D. Development of acceptance criteria for landfilling of waste: an approach based on impact modelling and scenario calculations. In: Proceedings SARDINIA-2001, Eighth International Waste Management and Landfill Symposium; S. Margherita di Pula, Cagliari (Italy); 2002. p. 711-721.

Hours M, Perrodin Y, al. Etude des polluants atmosphériques émis dans deux centres de stockage des ordures ménagères: caractérisation et mesure des niveaux d'exposition ; mise au point d'outils de suivi en vue de l'évaluation des risques sanitaires: Réseau Santé Déchets, Lyon; 2000. Rapport final d'étude.

INSERM. Dioxines dans l'environnement: quels risques pour la santé? Expertise collective de l'INSERM ; Paris; 2000.

Kjeldsen P, Christophersen M. Composition of leachate from old landfills in Denmark. Waste Manag.Res. 2001;19(3):249-256.

Kjeldsen P, Barlaz MA, Rooker AP, Baun A, Ledin A, Christensen TH. Present and long-term composition of MSW landfill leachate: a review. Crit.Rev.Environ.Sci.Technol.

2002;32(4):297-336.

Komilis DP, Stegmann R. The effect of landfill design and operation practices on waste degradation behaviour: a review. *Waste Manag. Res.* 1999;17:20-26.

Krug MN, Ham RK. Analysis of long-term leachate characteristics. In: *Proceedings Sardinia 1997, 6th International Landfill Symposium*; S. Margherita di Pula, Cagliari, Italy; 1997. p. 117-132.

Leduc R. *Guide de la modélisation de la dispersion atmosphérique*: Ministère de l'Environnement et de la Faune (Québec), Direction du milieu atmosphérique; 1998.

Lemière B, Seguin JJ, Le Guern C, Guyonnet D, Baranger P. *Guide sur le comportement des polluants dans le sol et les nappes*. Orléans: Editions BRGM; 2001. 119p.

Manca D, Birmingham B, Raha D. Toxicological screening of chemical emissions from municipal solid waste landfills. Application of a predictive frame work to a state-of-art facility. *Hum ecol Risk Assess* 1997;3(2):257-286.

McKone TE. Human exposure to volatile organic compounds in household tap water: the indoor inhalation pathway. *Environ.Sci.Technol.* 1987;21(12):1194-1201.

McKone T, Hall D, Kastenberg WE. *CalTOX version 2.3 Description of Modifications and Revisions*: University of California, Berkeley, Ernest Orlando Lawrence Berkeley National Laboratory; 1997.

Montgomery, Welkom. *Groundwater chemicals desk reference*. Lewis Publishers, Chelsea; 1990.

Nüsslein F, Feicht EA, Schulte-Hostede S, Kettrup A. Proposal on a standardized water risk assessment of toxic substances in soil. *Chemosphere* 1994;28(2):219-228.

OMS. *Guidelines for Air Quality*. WHO regional publications european series. Geneva: WHO (World Health Organization); 1999.

Parker T, Dottridge J, Kelly S. *Investigation of the composition and emissions of trace components in landfill gas*: Environmental Agency, Royaume Uni; 2002. R&D Technical Report P1-438/TR.

Paustenbach DJ, Rinehart WE, Sheehan PJ. The health hazards posed by chromium-contaminated soils in residential and industrial areas: conclusions of an expert panel. *Regulatory toxicology and pharmacology* 1991;13(2):195-222.

Ringard C, Rommens C, Duffa C. *Adaptation du programme de calcul du groupe Radioécologie Nord-Cotentin - Volet Radiologique au volet chimique*: IRSN; 2002. Note technique SEGR/SAER/01-64 Indice 1.

RIVM. *Variation in calculated exposure. Comparison of calculations with seven European exposure models.*; 2002. Report 711701030.

Robinson H, Knox K. *Pollution Inventory discharges to sewer or surface waters from landfill leachates*; 2001. REGCON 70.

Rollin AL, Jacquelin T. Geomembrane failures : lessons learned from geo-electrical leaks surveys: Leak Location Services inc., Canada; 1998.

Ruokojärvi P, Ettala M, Rahkonen P, Tarhanen J, Ruuskanen J. Polychlorinated dibenzo-P-dioxins and -furans (PCDDs and PCDFs) in municipal waste landfill fires. *Chemosphere* 1995;30(9):1697-1708.

SFSP (Société Française de Santé Publique). La pollution atmosphérique d'origine automobile et la santé publique: Collection Santé et Société n°4; 1996.

US-EPA (United States Environmental Protection Agency). Screening procedures for estimating the Air Quality Impact of Stationary Sources, revised: Office of Air and Radiation, Office of Air Quality Planning and Standards; October 1992. EPA-454/R-92-019.

US-EPA (United States Environmental Protection Agency). Screen3 Model User's guide; 1995. EPA-454/B-95-004.

US-EPA (United States Environmental Protection Agency). Exposure and Health assessment for 2,3,7,8-Tetrachlorodibenzo-p-dioxin (TCDD) and related compounds. Part 3 Integrated Summary and risk characterization for 2,3,7,8-Tetrachlorodibenzo-p-dioxin (TCDD) and related compounds; 2000.

Van Elsuve D, Fuvel V, Poulleau J. Analyses des rejets atmosphériques de torchères de biogaz: Ineris; 1999. Rapport Ineris DRC/MAPA/JPo21761n°99/16.

Zmirou D, Deloraine A, Saviuc P, Tillier C, Boucharlat A, Maury N. Short-term health effects of an industrial toxic waste landfill: a retrospective follow-up study in Montchanin, France. *Arch.Environ.Health* 1994;49(4):228-238.

CHAPITRE 4 RISQUES AIGUS, REPROTOXIQUES ET PERCEPTION D'ODEURS

ATSDR (Agency for Toxic Substances and Disease Registry). Toxicological Profiles for Trichloroethylene. In: US Department of Health and Human Services; 1997.

ATSDR (Agency for Toxic Substances and Disease Registry). Toxicological Profiles for Vinyl Chloride. In: US Department of Health and Human Services; 1997.

ATSDR (Agency for Toxic Substances and Disease Registry). Toxicological profile for Chlorinated Dibenzo-p-Dioxins. In: US Department of health and human services; 1998.

ATSDR (Agency for Toxic Substances and Disease Registry). Toxicological profile for Chloroethane. In: US Department of health and human services; 1998.

ATSDR (Agency for Toxic Substances and Disease Registry). Toxicological profile for Polychlorinated Biphenyls. In: US Department of health and human services; 2000.

ATSDR (Agency for Toxic Substances and Disease Registry). Toxicological profile for Aldrin/Dieldrin. In: US Department of health and human services; 2000.

ATSDR (Agency for Toxic Substances and Disease Registry). Toxicological profile for DDT, DDE and DDD. In: US Department of health and human services; 2000.

Devos M, Patte F, Rouault J, Laffort P. Standardized human olfactory threshold. Oxford University Press, Oxford; 1990.

Diosey P. Odor dispersion modeling : which models to use, how they work, what to expect. Air currents 1996;Section 4.

Dolk H, Vrijheid M, Armstrong B, Abramsky L, Bianchi F, Garne E et al. Risk of congenital anomalies near hazardous-waste landfill sites in Europe: the EUROHAZCON study. Lancet 1998;352(9126):423-427.

Dor F. Evaluation des risques sanitaires encourus par les populations riveraines après une exposition prolongée aux polluants de la décharge d'ordures ménagères au lieu dit La Bistade: InVS; Novembre 2000.

Elliott P, Briggs D, Morris S, de Hoogh C. Risk of adverse birth outcomes near landfill sites. Br. med. J. 2001a;323(7325):363-368.

Elliott P, Morris S, Briggs D, de Hoogh C, Hurt C, Kold Jensen T et al. Birth outcomes and selected cancers in populations living near landfill sites; 2001b August 2001. Report to the department of Health.

Elliott P, Jarup L, Briggs D, de Hoogh C, Morris S, Hurt C et al. Cancer risks in populations living near landfill sites in Great Britain. Br. J. Cancer 2002;86(11):1732-1736.

ENSP. Les risques non microbiologiques associés au compostage: Etude FNADE-MEDD réalisée par l'ENSP; Août 2002. 142p.

Gingras B, Guy C, Pagé T. Chapitre 19 : Odeurs. In: TEC&DOC, editor. Environnement et Santé Publique. Fondements et pratiques; 2003. 1023p.

Goldberg MS, Siemiatycki J, Dewar R, Desy M, Riberdy H. Risks of developing cancer relative to living near a municipal solid waste landfill site in Montreal, Quebec, Canada. Arch. environ. Health 1999;54(4):291-296.

INRS. Produits chimiques cancérigènes, mutagènes, toxiques pour la reproduction - classification réglementaire. Cahiers de notes documentaires - Hygiène et sécurité du travail 2ème trimestre 2002(N°187).

Johnson BL. A review of the effects of hazardous waste on reproductive health. Am. J. Obstetrics Gynecol. 1999;181(1):S12-S16.

Leduc R. Guide de la modélisation de la dispersion atmosphérique: Ministère de l'Environnement et de la Faune (Québec), Direction du milieu atmosphérique; 1998.

Pagé T, Guy C. Odor dispersion modeling. In: Air & Waste Management Association's 90th Annual Meeting & Exhibition; 8-13 Juin 1997; Toronto, Ontario, Canada; 1997.

Pope RJ, Diosey P. Odor dispersion : models and methods. CLEARWATERS 2000;30(2).

Rousselin X, Bosio E, Falcy M. Comparaison des seuils olfactifs de substances chimiques

avec des indicateurs de sécurité utilisés en milieu professionnel. Cahier de notes documentaires INRS 1994(156):315-327.

Senante E, Galtier L, Békaert C, Lambolez-Michel L, Budka A. Odours management at MSW landfill sites : odours sources, odorous compounds and control measures. In: CISA publisher, editor. Proceedings Sardinia 2003, Ninth International Waste Management and Landfill Symposium; to be published; Cagliari; 2003.

Shusterman D. Critical review : the health significance of environmental odor pollution. Arch.Environ.Health 1992;47:76-87.

Termonia A, Termonia M. Characterisation and on-site monitoring of odorous organic compounds in the environment of a landfill site. Intern. J. Environ. Anal. Chem. 1998;73(1):43-57.

US-EPA (United States Environmental Protection Agency). Screen3 Model User's guide; 1995. EPA-454/B-95-004.

US-EPA (United States Environmental Protection Agency). Guidelines for developmental toxicity risk assessment; décembre 1991. EPA/600/FR-91/001.

US-EPA (United States Environmental Protection Agency). Phenol, CASRN 108-95-2. In: IRIS (Integrated Risk Information System); 2003.

Vrijheid M, Dolk H, Armstrong B, Abramsky L, Bianchi F, Fazarinc I et al. Chromosomal congenital anomalies and residence near hazardous waste landfill sites. Lancet 2002;359(9303):320-322.